


Springer fibers and Richardson varieties

Slides available at snkarp.github.io



Steven N. Karp (University of Notre Dame)
joint work with Martha Precup

April 14th, 2025
Georgia Institute of Technology

Flag variety $\text{Fl}_n(\mathbb{C})$

- The (*complete*) *flag variety* $\text{Fl}_n(\mathbb{C})$ is the set of all flags $F_\bullet$ in $\mathbb{C}^n$:

$$F_\bullet = (0 = F_0 \subset F_1 \subset \cdots \subset F_{n-1} \subset F_n = \mathbb{C}^n), \quad \dim(F_j) = j \text{ for all } j.$$

Equivalently, $\text{Fl}_n(\mathbb{C}) = \text{GL}_n(\mathbb{C})/B_+$.

- e.g. $n = 3$

$$F_\bullet = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \\ 3 & 0 & 0 \end{bmatrix} \iff F_0 = 0, \quad F_1 = \left\langle \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix} \right\rangle, \quad F_2 = \left\langle \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\rangle, \quad F_3 = \mathbb{C}^3$$

- An $n \times n$ nilpotent matrix M a flag $F_\bullet$ are *compatible* if

$$M(F_j) \subseteq F_{j-1} \quad \text{for all } j \geq 1.$$

- e.g. If $M := \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ then M and $F_\bullet$ are compatible.

Springer fiber $\mathcal{B}_\lambda$ (1969)

- Let λ be a *partition* of n (i.e. a weakly decreasing sequence of positive integers summing to n). Let M_λ be the $n \times n$ nilpotent matrix of type λ in Jordan form. Every nilpotent matrix is conjugate to a unique M_λ .

- e.g. $M_{(3)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, $M_{(1,1,1)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $M_{(2,1)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

- The *Springer fiber* $\mathcal{B}_\lambda \subseteq \text{Fl}_n(\mathbb{C})$ is the set of flags compatible with M_λ .

- e.g. $\mathcal{B}_{(3)} = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$, $\mathcal{B}_{(1,1,1)} = \text{Fl}_3(\mathbb{C})$,

$$\mathcal{B}_{(2,1)} = \left\{ \begin{bmatrix} a & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \right\} \sqcup \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & b & 1 \\ 0 & 1 & 0 \end{bmatrix} \right\} \sqcup \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$$

- The dimension of $\mathcal{B}_\lambda$ is $\sum_{i \geq 1} (i-1)\lambda_i$.

Why study Springer fibers?

- $\mathcal{B}_\lambda$ is a fiber in the *Springer resolution* of the nilpotent cone of $\mathfrak{sl}_n(\mathbb{C})$.
- Springer (1976): The cohomology $H^*(\mathcal{B}_\lambda)$ is a representation of the symmetric group S_n of all permutations of $\{1, \dots, n\}$. Moreover, $H^{\text{top}}(\mathcal{B}_\lambda)$ is the irreducible Specht module indexed by λ .
- Springer fibers and generalizations (such as Hessenberg varieties) are related to chromatic quasisymmetric functions and Macdonald polynomials.
- The geometry of irreducible components is related to the combinatorics of Catalan numbers, webs, and standard tableaux.
- A *standard (Young) tableau* σ of shape λ is a filling of the diagram of λ with $1, \dots, n$ (each used once) which is increasing along rows and columns.

• e.g. $\lambda = (4, 2, 2) =$

$\sigma =$

1	3	5	6
2	4		
7	8		

$\sigma[3] =$

1	3
2	

- For $1 \leq j \leq n$, let $\sigma[j]$ denote the tableau formed by entries $1, \dots, j$ of σ .

Irreducible components $\mathcal{B}_\sigma$ of Springer fibers

- Given $F_\bullet \in \mathcal{B}_\lambda$, let $\text{JF}(F_\bullet)$ denote the standard tableau σ such that for all j , the shape of $\sigma[j]$ is the Jordan type of M_λ acting on $\mathbb{C}^n/F_{n-j}$.

- e.g. $\lambda = (2, 1)$, $M_\lambda = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $F_\bullet = \begin{bmatrix} 1 & 0 & 0 \\ 0 & b & 1 \\ 0 & 1 & 0 \end{bmatrix}$

$$M_\lambda = 0 \text{ on } \mathbb{C}^3/F_1 \Rightarrow \text{shape}(\sigma[2]) = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \Rightarrow \text{JF}(F_\bullet) = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}$$

- Spaltenstein (1976): The irreducible components of $\mathcal{B}_\lambda$ are precisely

$$\mathcal{B}_\sigma := \overline{\mathcal{B}_\sigma^\circ}, \quad \text{where } \mathcal{B}_\sigma^\circ := \{F_\bullet \in \mathcal{B}_\lambda : \text{JF}(F_\bullet) = \sigma\}.$$

- e.g. $\mathcal{B}_{\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}}^\circ = \left\{ \begin{bmatrix} a & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \right\}, \quad \mathcal{B}_{\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}}^\circ = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & b & 1 \\ 0 & 1 & 0 \end{bmatrix} \right\} \sqcup \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$

Evacuation on standard tableaux

- Let $\sigma \mapsto \sigma^\vee$ denote *evacuation* (or the *Schützenberger involution*) on standard tableaux. We obtain $\sigma^\vee$ from σ by repeating these steps:
 - delete the box in the top-left corner;
 - slide the empty box to the southeast boundary (via *jeu-de-taquin*);
 - fill the empty box with the largest unused number (starting with n) and freeze it.

e.g. $\sigma =$

1	3	5	6
2	4		
7	8		

 $\rightsquigarrow$ $\sigma^\vee =$

1	2	5	7
3	4		
6	8		

- van Leeuwen (2000): If $F_\bullet \in \mathcal{B}_\sigma^\circ$, then the Jordan type of M_λ acting on F_j is the shape of $\sigma^\vee[j]$, for all j .
- We can similarly recover the *Robinson–Schensted correspondence*.

Total positivity

- *Totally positive matrices* (matrices whose minors are all positive) have been studied since the 1930's. Gantmakher–Krein (1937) showed that square totally positive matrices have positive eigenvalues.

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \quad \begin{aligned} \lambda_1 &= 10.6031 \dots \\ \lambda_2 &= 1.2454 \dots \\ \lambda_3 &= 0.1514 \dots \end{aligned}$$

- Lusztig (1994) introduced total positivity for algebraic groups G and flag varieties G/P . It is connected to representation theory, cluster algebras, electrical networks, combinatorics, topology, Teichmüller theory, tropical and real algebraic geometry, scattering amplitudes, knot theory, ...
- The *totally nonnegative flag variety* $\mathrm{Fl}_n^{\geq 0}$ is the set of flags which have a matrix representative whose left-justified minors are all nonnegative.

- e.g. $F_\bullet = \begin{bmatrix} \boxed{2} & \boxed{-3} & 1 \\ 4 & -1 & 0 \\ \boxed{1} & \boxed{0} & 0 \end{bmatrix} \in \mathrm{Fl}_3^{\geq 0}$ **minor = 3**

Cell decomposition of $Fl_n^{\geq 0}$




- Lusztig (1994), Rietsch (1999): $Fl_n^{\geq 0}$ has a cell decomposition

$$Fl_n^{\geq 0} = \bigsqcup_{v \leq w \text{ in } S_n} R_{v,w}^{\geq 0},$$

where $R_{v,w}^{\geq 0}$ is the totally positive part of the *Richardson variety*

$$R_{v,w} := \overline{B_- v B_+ / B_+} \cap \overline{B_+ w B_+ / B_+}.$$

- Galashin–Karp–Lam (2022): $Fl_n^{\geq 0}$ is a regular CW complex.
- e.g. $n = 3$




Totally nonnegative Springer fiber $\mathcal{B}_\lambda^{\geq 0}$

- Lusztig (2020): The *totally nonnegative Springer fiber* is a subcomplex of $\mathrm{Fl}_n^{\geq 0}$:

$$\mathcal{B}_\lambda^{\geq 0} := \mathcal{B}_\lambda \cap \mathrm{Fl}_n^{\geq 0} = \bigsqcup_{\substack{v \leq w \text{ in } S_n, \\ R_{v,w} \subseteq \mathcal{B}_\lambda}} R_{v,w}^{\geq 0}.$$

Its top-dimensional cells are indexed by $R_{v,w}$'s which equal some $\mathcal{B}_\sigma$.

- e.g.



Richardson envelope of $\mathcal{B}_\sigma$

Problem

When is the irreducible component $\mathcal{B}_\sigma$ equal to a Richardson variety $R_{v,w}$?
That is, describe the top-dimensional cells of $\mathcal{B}_\lambda^{\geq 0}$ in terms of tableaux.

- Additional motivation: much more is known about $R_{v,w}$ than $\mathcal{B}_\sigma$.

Theorem (Karp, Precup (2025+))

There exists a unique minimal Richardson variety R_{v_σ, w_σ} containing $\mathcal{B}_\sigma$.

- e.g.

$$\mathcal{B}_{\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}} = \overline{\left\{ \begin{bmatrix} a & \boxed{1} & 0 \\ 0 & 0 & \boxed{1} \\ \boxed{1} & 0 & 0 \end{bmatrix} \right\}} = \overline{\left\{ \begin{bmatrix} \boxed{1} & 0 & 0 \\ 0 & 0 & \boxed{1} \\ \frac{1}{a} & \boxed{1} & 0 \end{bmatrix} \right\}} = R_{132, 312}$$

$$\mathcal{B}_{\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array}} = \overline{\left\{ \begin{bmatrix} a & b & \boxed{1} & 0 \\ 0 & a & 0 & \boxed{1} \\ \boxed{1} & 0 & 0 & 0 \\ 0 & \boxed{1} & 0 & 0 \end{bmatrix} \right\}} = \overline{\left\{ \begin{bmatrix} \boxed{1} & 0 & 0 & 0 \\ 0 & \boxed{1} & 0 & 0 \\ \frac{1}{a} & -\frac{b}{a^2} & \boxed{1} & 0 \\ 0 & \frac{1}{a} & 0 & \boxed{1} \end{bmatrix} \right\}} \subsetneq R_{1234, 3412}$$

Reading words of σ

- We can find the Richardson envelope R_{v_σ, w_σ} of $\mathcal{B}_\sigma$ as follows:
 - v_σ^{-1} is the *top-down reading word* of $\sigma^\vee$; and
 - $w_0 w_\sigma^{-1} w_0$ is the *reading word* of σ , where $w_0 := (j \mapsto n+1-j) \in S_n$.

• e.g.

$$\sigma =$$

1	3	5	6
2	4		
7	8		

$\longleftrightarrow$

$$\sigma^\vee =$$

1	2	5	7
3	4		
6	8		

$$v_\sigma^{-1} = 12573468$$

$\Rightarrow$

$$v_\sigma = 12563748$$

$$w_0 w_\sigma^{-1} w_0 = 78241356$$

$\Rightarrow$

$$w_\sigma = 78125364$$

$$w_0 = 87654321$$

- Pagnon (2003), Pagnon–Ressayre (2006): X_{w_σ} is the minimal Schubert variety containing $\mathcal{B}_\sigma$.

Richardson tableaux

- A *Richardson tableau* is a standard tableau σ such that for all $1 \leq j \leq n$, if j appears in row r of σ , then either $r = 1$ or the **largest entry** of $\sigma[j-1]$ in row $r-1$ is greater than every entry in rows $\geq r$.

- e.g.

$$\sigma = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 4 & 6 \\ \hline 2 & 7 & & \\ \hline 5 & 8 & & \\ \hline \end{array}$$

Richardson

$$\tau = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 5 & 6 \\ \hline 2 & 4 & & \\ \hline 7 & 8 & & \\ \hline \end{array}$$

not Richardson

- All hook-shaped tableaux are Richardson. A two-rowed tableau is Richardson if and only if its second row has no two consecutive entries.

Theorem (Karp, Precup (2025+))

Let σ be a standard tableau of shape λ . The following are equivalent:


- (i) $\mathcal{B}_\sigma$ is equal to some Richardson variety $R_{v,w}$;
- (ii) $\mathcal{B}_\sigma = R_{v_\sigma, w_\sigma}$; and
- (iii) σ is a Richardson tableau.

Slide characterization of Richardson tableaux

Theorem (Karp, Precup (2025+))

A standard tableau σ is Richardson if and only if every evacuation slide of σ (used to calculate $\sigma^\vee$) is L-shaped.

• e.g.



• Important (non-obvious) fact: if σ is Richardson, then so is $\sigma^\vee$.

Enumeration of Richardson tableaux of fixed shape

Theorem (Karp, Precup (2025+))

The number of Richardson tableaux of shape $\lambda = (\lambda_1, \dots, \lambda_\ell)$ is

$$\binom{\lambda_{\ell-1}}{\lambda_\ell} \binom{\lambda_{\ell-2} + \lambda_\ell}{\lambda_{\ell-1} + \lambda_\ell} \binom{\lambda_{\ell-3} + \lambda_{\ell-1} + \lambda_\ell}{\lambda_{\ell-2} + \lambda_{\ell-1} + \lambda_\ell} \cdots \binom{\lambda_1 + \lambda_3 + \lambda_4 + \cdots + \lambda_\ell}{\lambda_2 + \lambda_3 + \lambda_4 + \cdots + \lambda_\ell}.$$

- e.g. $\lambda = (3, 2, 1) \rightsquigarrow \binom{2}{1} \binom{3+1}{2+1} = 2 \cdot 4 = 8$

1	2	4
3	5	
6		

1	2	5
3	6	
4		

1	3	4
2	5	
6		

1	3	5
2	4	
6		

1	3	5
2	6	
4		

1	3	6
2	4	
5		

1	4	5
2	6	
3		

1	4	6
2	5	
3		











- A q -analogue holds using the major index on standard tableaux.

Enumeration of Richardson tableaux of fixed size

Theorem (Karp, Precup (2025+))

The number of Richardson tableaux of size n is the Motzkin number M_n .

- e.g. $n = 4$, $M_4 = 9$



Singular locus and cohomology class of $\mathcal{B}_\sigma$

- There is no known general algorithm to find the singular locus of $\mathcal{B}_\sigma$.
- Billey–Coskun (2012) explicitly describe the singular locus of $R_{v,w}$. Thus we can calculate the singular locus of $\mathcal{B}_\sigma$ if σ is a Richardson tableau.

Problem

For which Richardson tableaux σ is $\mathcal{B}_\sigma$ smooth?

- In all examples of Richardson tableaux σ that we checked (including all such σ with at most two columns or at most three rows), $\mathcal{B}_\sigma$ is smooth.

Problem

Expand the cohomology class $[\mathcal{B}_\sigma]$ in the Schubert basis of $H^(\mathrm{Fl}_n(\mathbb{C}))$.*

- If σ is Richardson, this is equivalent to expanding $\mathfrak{S}_{v_\sigma} \mathfrak{S}_{w_0 w_\sigma}$ in the basis of Schubert polynomials, which was solved by Spink–Tewari (2025+).

Thank you!